

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.Sciencedirect.com)

## International Journal of Surgery

journal homepage: [www.theijs.com](http://www.theijs.com)

## Original research

Concurrent rib and pelvic fractures as an indicator of solid abdominal organ injury<sup>☆</sup>

Ammar Al-Hassani<sup>a,\*</sup>, Ibrahim Afifi<sup>a</sup>, Husham Abdelrahman<sup>a</sup>, Ayman El-Menyar<sup>b,c</sup>,  
Ammar Almadani<sup>a</sup>, Jan Recicar<sup>a</sup>, Hassan Al-Thani<sup>a</sup>, Kimball Maull<sup>a,d</sup>, Rifat Latifi<sup>a,b,e</sup>

<sup>a</sup> Section of Trauma Surgery, Hamad General Hospital, Qatar<sup>b</sup> Clinical Medicine, Weill Cornell Medical College, Doha, Qatar<sup>c</sup> Clinical Research, Trauma Surgery Section, Hamad General Hospital, Qatar<sup>d</sup> University of Pittsburgh Medical Center, USA<sup>e</sup> Department of Surgery, University of Arizona, Tucson, AZ, USA

## ARTICLE INFO

## Article history:

Received 14 September 2012

Received in revised form

16 March 2013

Accepted 8 April 2013

Available online 17 April 2013

## Keywords:

Blunt trauma

Multiple rib fracture

Solid organ injury

Pelvic fracture

## ABSTRACT

**Objectives:** To study the association of solid organ injuries (SOIs) in patients with concurrent rib and pelvic fractures.

**Methods:** Retrospective analysis of prospectively collected data from November 2007 to May 2010. Patients' demographics, mechanism of injury, Injury severity scoring, pelvic fracture, and SOIs were analyzed. Patients with SOIs were compared in rib fractures with and without pelvic fracture.

**Results:** The study included 829 patients (460 with rib fractures  $\pm$  pelvic fracture and 369 with pelvic fracture alone) with mean age of  $35 \pm 12.7$  years. Motor vehicle crashes (45%) and falls from height (30%) were the most common mechanism of injury. The overall incidence of SOIs in this study was 22% (185/829). Further, 15% of patient with rib fractures had associated pelvic fracture. SOI was predominant in patients with concurrent rib fracture and pelvic fracture compared to ribs or pelvic fractures alone (42% vs. 26% vs. 15%, respectively,  $p = 0.02$ ).

**Conclusions:** Concurrent multiple rib fractures and pelvic fracture increases the risk of SOI compared to either group alone. Lower RFs and pelvic fracture had higher association for SOI and could be used as an early indicator of the presence of SOIs.

© 2013 Surgical Associates Ltd. Published by Elsevier Ltd. All rights reserved.

## 1. Introduction

Rib fractures are common chest injuries resulting from high force blunt trauma. It is difficult to determine the exact incidence of rib fractures among seriously injured patients due to insensitivity of routine chest radiography.<sup>1</sup> Furthermore, the clinical bedside diagnosis of intraabdominal injuries is difficult in those patients who are at risk for intraabdominal injury.<sup>2,3</sup> At present, accurate diagnosis of intra-abdominal injuries in hemodynamically stable patients is primarily based on abdominal CT scanning.<sup>4,5</sup>

<sup>☆</sup> This study was presented in part in Trauma Association of Canada Annual Scientific Meeting, Banff, April 07–08, 2011.

\* Corresponding author. Section of Trauma Surgery, Department of Surgery, Hamad General Hospital, P.O. Box 3050, Doha, Qatar. Tel.: +974 4439 6152 (Office).

E-mail addresses: [TraumaResearch@hmc.org.qa](mailto:TraumaResearch@hmc.org.qa), [ammam\\_alhassani@yahoo.com](mailto:ammam_alhassani@yahoo.com) (A. Al-Hassani).

The majority of the patients with rib fracture due to blunt trauma have significant associated injuries.<sup>6–9</sup>

Patient with lower rib fractures of the right side are more likely to have liver injury, whereas left-sided fractures are more susceptible toward splenic injury.<sup>9,10</sup> Pelvic fracture is an indicator of severe trauma and is often associated with major intraabdominal injuries.<sup>11</sup> Though the incidence of pelvic fractures after blunt trauma is relatively high, it had lesser association with severe complications.<sup>11</sup> Eastridge et al., showed that not only the pelvic fracture but also, the associated complications such as neurologic (27%), thoracic (26%) and abdominal (14%) injuries correlated with the collision force and severity in blunt trauma patients.<sup>12</sup> It is also noted that in patients with severe pelvic fractures (Abbreviated Injury Score  $\geq 4$ ), the incidence of associated injuries was notably higher (31%).<sup>11</sup>

We, previously, reported on association of lower rib fractures with an increased incidence of solid organ injury (SOI).<sup>13</sup> To the best

of our knowledge, there is no report available on the association between rib, pelvic fracture and abdominal SOI.

Using rib and pelvic fractures as an indicator of blunt trauma force, the association of both fractures with intra-abdominal injuries was studied with the objective to determine whether a significant relationship exists between the occurrence of these fractures and the incidence of associated SOI.

## 2. Methods

Hamad General Hospital, the tertiary care medical facility in Doha, Qatar, provides all levels of medical services to the entire population and receives all major trauma patients from the country. Trauma surgeons from initial assessment and resuscitation, thru operation, intensive care and in-hospital care, manage the injured patients. Following an initial review of our experience, data were collected and analyzed retrospectively for all patients admitted with multiple rib fractures and presence of pelvic fracture during the period from November 2007 to May 2010.

Demographics, mechanism of injury, Injury Severity Score (ISS), presence of pelvic fracture, rib fracture (number and location) and associated abdominal SOIs were analyzed for all cases and controls. Initial clinical assessment was performed according to Advanced Trauma Life Support protocols. All patients included in the study were assessed by initial supine chest and pelvic radiographs followed by computed tomography scan of chest and abdomen. The solid organs injury (SOI) of the abdomen constituted liver, spleen, and kidneys. SOI was compared in rib fracture patients with and without pelvic fractures. Data were presented as proportions, mean  $\pm$  standard deviation (SD) and range as appropriate. Baseline demographic characteristics, presentation and outcomes were compared using the student-*t* test for continuous variables and Pearson chi-square ( $\chi^2$ ) test for categorical variables. A significant difference was considered when the *P* value was less than 0.05. All data analyses were carried out using the Statistical Package for Social Sciences version 18 (SPSS Inc. USA). The study was approved by Medical Research Center, HMC, Qatar (IRB#10062/10).

## 3. Results

The study cohort included 829 patients who sustained blunt traumatic injury, of which 460 had multiple rib fractures (with and without associated pelvic fracture) and 369 had isolated pelvic fracture. The majority of patients were males (93%), with a mean age of  $35 \pm 12.7$  years.

The mechanism of injury: Motor vehicle crash (206; 45%) and fall from height (139; 30%) were the most common mechanism of injury (Fig. 1).

Fig. 2 shows the overview of the study results. In the study group, concurrent rib and pelvic fracture was observed in 69 (15%) cases, whereas, 391 (85%) cases had rib fracture alone. Moreover, 369 patients with pelvic fracture alone were included as controls. The overall incidence of SOIs in this study was 22% (185/829). SOIs were identified in 130 (28.3%) patients in the study group and in 55 (15%) patients of the control group. Fig. 3 shows that SOI was predominant in patients with concurrent rib fracture and pelvic fracture compared to ribs or pelvic fractures alone (42% vs. 26% vs. 15%, respectively,  $p = 0.02$ ).




Fig. 1. Mechanism of injury.

The most common injured organ was spleen followed by liver and kidney. Median abdomen AIS was 3 (1–4) whereas median ISS was 13 (1–45). ISS was higher in the study group when compared to the control group [(13 (4–66) vs. 9 (4–50);  $p < 0.001$ )]. Further, in patients with SOI, the presence of pelvic fractures was associated with higher ISS in comparison to those who had isolated rib fracture (Table 1).

SOIs were graded according to American Association for the Surgery of Trauma – Organ Injury Scale (AAST/OIS) as mild to moderate (grade I, II) in 123 patients and severe (grade III, IV) in 51 patients. No association was observed between pelvic fracture and number of rib fractures, as the majority (78%) of cases had  $<5$  fractured ribs.

Table 2 shows the number of patients presenting with concurrent pelvic fracture and rib fracture (either isolated or overlapping rib zones). The majority of patients with pelvic fractures were reported in the presence of overlapping middle and upper rib zones (42%) followed by overlapping lower and middle rib zones (26%).

Table 3 shows the number of patients presenting with solid organ injury in the presence of concurrent pelvic fracture and rib fracture (either isolated or overlapping rib zones). SOIs were mainly encountered in patients with pelvic fracture in the presence of overlapping lower and middle rib zones (62%) followed by isolated lower rib zone (50%).

## 4. Discussion

The present study reports that the presence of pelvic fracture increases the incidence of associated SOI in patients with rib fracture. The overall incidence of SOIs in this study was 22%. The incidence of SOI was significantly higher in patients with combined rib and pelvic fractures compared to either rib or pelvic fractures alone.

An increased risk of abdominal SOIs in patients of lower rib fracture has been reported earlier.<sup>14–16</sup> Consistent with these reports, this study also shows higher association of SOI with lower rib fractures and overlapping lower and middle rib zones which also corroborates with our previous report.<sup>13</sup>

The rising incidence of road traffic crashes is a most important public health problem in the civil society. In particular, blunt chest injury and pelvic fractures are associated with increased morbidity and mortality. The chest wall and soft tissues are the most commonly affected sites by blunt trauma. The site of fracture has important impact on the clinical presentation. Fractured ribs 4–10 are more frequent in trauma, whereas, fractured ribs 8–12 increases the likelihood of the presence of associated abdominal injuries.<sup>14</sup>

Surprisingly, there are only few reports on the potential additive impact of pelvic or long bone fractures on the likelihood of associated abdominal injuries. Shweiki et al. reviewed 476 hospitalized traumatic rib fracture patients and concluded that in patients with rib fracture, presence of pelvic fractures and long bone fractures did not increase the likelihood of associated SOI.<sup>9</sup> In contrast, our study showed increased association of SOI and concurrent multiple rib and pelvic fractures. It also demonstrates high predictability for SOI with concurrent pelvic and lower rib fractures. Pelvic fractures are reportedly the third most common cause of death in motor vehicle crashes, after central nervous system and chest injuries.<sup>17</sup> Pelvic injuries, especially those involving disruption of the pelvic ring, signify high energy trauma. Such injuries are seen in motor vehicle crashes and in falls from considerable height, precisely the largest groups represented in this study.

Parriera et al., demonstrated that outcome was more closely related to the associated injuries than to the pelvic injuries per se.<sup>18</sup> Demetriades et al., also showed high incidence of associated




Fig. 2. Overview of study results.




Fig. 3. Incidence of solid organ injuries in relation to pelvic and rib fracture.

abdominal injuries among adults and pediatric trauma patients and liver was the most common injured organ.<sup>11,19</sup> In patients with complex pelvic fractures, the spleen was found to be the most frequently injured solid organ followed by liver.<sup>20,21</sup> In our study, the incidence of liver and splenic injuries were comparable in patients with concurrent rib and pelvic fractures.

The association of intra-abdominal injuries with pelvic fractures is well recognized. Bond et al., reported an incidence of 20% of associated intra-abdominal injuries in pediatric pelvic fractures patients.<sup>22</sup> The high incidence of associated abdominal injuries (42%) in this study was probably due to exclusion of simple pelvic avulsion fractures and higher median Injury Severity Score.

Table 1

Injury severity score according to rib and pelvic fracture with associated solid organ injury.

Solid organ injury	Injury severity score (median; range)
Rib + pelvic fracture (n = 29)	22 (9–41)
Rib fracture alone (n = 101)	13 (5–45)
Pelvic fracture alone (n = 55)	22 (8–50)
No solid organ injury (n = 314)	13 (4–60)

Table 2

Number of patients presenting with concurrent pelvic fracture and rib fracture (either isolated or overlapping rib zones).

	Lower ribs	Middle ribs	Upper ribs
Lower ribs	8	26	—
Middle ribs	26	9	42
Upper ribs	—	42	18

Table 3

Number of patients presenting with solid organ injury in the presence of concurrent pelvic fracture and rib fracture (either isolated or overlapping rib zones).

	Lower ribs	Middle ribs	Upper ribs
Lower ribs	4/8 (50%)	16/26 (62%)	—
Middle ribs	16/26 (62%)	4/9 (44%)	18/42 (43%)
Upper ribs	—	18/42 (43%)	6/18 (33%)

#### 4.1. Limitations

The important limitations of the study included the retrospective nature of the study, small number of patients and the overlap between the locations of fractured ribs.

Particularly, there was an overlap between the adjacent lower and middle ribs and also between middle and upper ribs. Further, the types of pelvic fracture were not given clearly.

#### 5. Conclusions

The concurrent clinical findings of fractures of the lower ribs and, pelvic fracture after blunt trauma pose a high risk for intra-abdominal SOI. Association of multiple rib fractures and pelvic fracture increases the risk of SOI compared to either group occurring separately.

Contrarily, rib fracture patients without concurrent pelvic fracture are at a lower risk of intra-abdominal injury. Therefore, the presence of pelvic fracture and multiple ribs fracture should be considered as a potential indicator for possible solid organ injury.

#### Ethical approval

The study was approved by Medical Research Center, Hamad Medical Corporation, Qatar (IRB#10062/10).

**Funding**

None.

**Author contribution**

Ammar Al-Hassani (Lead investigator & writing manuscript), Ibrahim Afifi (data collection), H Abdelrahman (data collection & review manuscript), Ammar Al Madani (data collection), J Recicar (review manuscript), Ayman El-Menyar (data analysis & writing & review manuscript), Rifat Latifi (review manuscript), Hassan Al Thani (review manuscript), K Maull (review manuscript).

**Conflict of interest**

None.

**Acknowledgment**

We thank all the trauma surgery staff for their cooperation. The authors have no conflict of interest and no financial issues to disclose. All authors read and approved the manuscript.

**References**

- Exadaktylos AK, Sclabas G, Schmid SW, Schaller B, Zimmermann H. Do we really need routine computed tomographic scanning in the primary evaluation of blunt chest trauma in patients with "normal" chest radiograph? *J Trauma* 2001 Dec;**51**(6):1173–6.
- Poletti PA, Mirvis SE, Shanmuganathan K, Takada T, Killeen KL, Perlmutter D, et al. Blunt abdominal trauma patients: can organ injury be excluded without performing computed tomography? *J Trauma* 2004;**57**:1072–81.
- Schurink GW, Bode PJ, van Luijt PA, van Vugt AB. The value of physical examination in the diagnosis of patients with blunt abdominal trauma: a retrospective study. *Injury* 1997;**28**:261–5.
- Stuhlfaut JW, Anderson SW, Soto JA. Blunt abdominal trauma: current imaging techniques and CT findings in patients with solid organ, bowel, and mesenteric injury. *Semin Ultrasound CT MR* 2007;**28**:115–29.
- Radwan MM, Abu-Zidan FM. Focussed assessment sonograph trauma (FAST) and CT scan in blunt abdominal trauma: surgeon's perspective. *Afr Health Sci* 2006;**6**(3):187–90.
- Shorr RM, Crittenden M, Indeck M, Hartunian SL, Rodriguez A. Blunt thoracic trauma. Analysis of 515 patients. *Ann Surg* 1987;**206**(2):200–5.
- Ziegler DW, Agarwal NN. The morbidity and mortality of rib fractures. *J Trauma* 1994;**37**(6):975–9.
- Holcomb JB, McMullin NR, Kozar RA, Lygas MH, Moore FA. Morbidity from rib fractures increases after age 45. *J Am Coll Surg* 2003 Apr;**196**(4):549–55.
- Shweiki E, Klena J, Wood GC, Indeck M. Assessing the true risk of abdominal solid organ injury in hospitalized rib fracture patients. *J Trauma* 2001 Apr;**50**(4):684–8.
- Yoganandan N, Pintar FA, Gennarelli TA, Maltese MR. Patterns of abdominal injuries in frontal and side impacts. *Annu Proc Assoc Adv Automot Med* 2000;**44**:17–36.
- Demetriades D, Karaiskakis M, Toutouzas K, Alo K, Velmahos G, Chan L. Pelvic fractures: epidemiology and predictors of associated abdominal injuries and outcomes. *J Am Coll Surg* 2002 Jul;**195**(1):1–10.
- Eastridge BJ, Burgess AR. Pedestrian pelvic fractures: 5-year experience of a major urban trauma center. *J Trauma* 1997 Apr;**42**(4):695–700.
- Al-Hassani A, Abdulrahman H, Afifi I. Rib fracture patterns predict thoracic chestwall and abdominal solid organ injury. *Am Surg* 2010 August;**76**(8):888–91 (4).
- Lee J, Harris Jr JH, Duke Jr JH, Williams JS. Noncorrelation between thoracic skeletal injuries and acute traumatic aortic tear. *J Trauma* 1997;**43**(3):400–4.
- Poole Jr GV, Myers RT. Morbidity and mortality rates in major blunt trauma to the upper chest. *Ann Surg* 1981;**193**:70–5.
- Lee RB, Bass SM, Morris Jr JA, MacKenzie EJ. Three or more rib fractures as an indicator for transfer to a level I trauma center: a population-based study. *J Trauma* 1990;**30**:689–94.
- Giannoudis PV, Grotz MR, Tzioupis C, Dinopoulos H, Wells GE, Bouamra O, et al. Prevalence of pelvic fractures, associated injuries, and mortality: the United Kingdom perspective. *J Trauma* 2007;**63**(4):875–83.
- Parreira JG, Coimbra R, Rasslan S, Oliveira A, Fregoneze M, Mercadante M. The role of associated injuries on outcome of blunt trauma patients sustaining pelvic fractures. *Injury* 2000;**31**:677–82.
- Demetriades D, Karaiskakis M, Velmahos G, Alo K, Murray J, Chan L. Pelvic fractures in pediatric and adult trauma patients: are they different injuries? *J Trauma-injury Infect Crit Care* 2003 June;**54**(6):1146–51.
- Paienda GS, Seitz H, Mousavi M, Vecsei V. Concomitant intra-abdominal injuries in pelvic trauma. *Wien Klin Wochenschr* 1998 Dec;**110**(23):834–40.
- Lunsjo K, Tados A, Hauggaard A, Blomgren R, Kopke J, Abu-Zidan A. Associated injuries and not fracture instability predict mortality in pelvic fractures: a prospective study of 100 patients. *J Trauma* 2007;**62**(3):687.
- Bond SJ, Gotschall CS, Eichelberger MR. Predictors of abdominal injury in children with pelvic fractures. *J Trauma* 1991;**31**:1169–72.